

Bibliografia

Direct anterior approach versus posterolateral approach in total hip arthroplasty: effects on early post-operative rehabilitation period. Faldini C, Perna F, Mazzotti A, Stefanini N, Panciera A, Geraci G, Mora P, Traina F. *J Biol Regul Homeost Agents*. 2017 Oct-Dec;31(4 suppl 1):75-81

Modified minimally invasive direct anterior approach through a bikini incision for total hip arthroplasty: technique and results in young female patients. Faldini C, Mazzotti A, Perna F, Stefanini N, Panciera A, Pilla F, Ruffilli A, Neonakis EM, Traina F. *J Biol Regul Homeost Agents*. 2017 Oct-Dec;31(4 suppl 1):83-89.

Anterolateral soft tissue sparing approach to the hip. Surgical technique. Toni A, Traina F, De Fine M, Garagnani L, Cervini A, Tassinari E. *Chir Organi Mov*. 2008 May;92(1):23-7. doi: 10.1007/s12306-008-0040-9. Epub 2008 Apr 12.

Outcomes of total hip replacement in patients with slipped capital femoral epiphysis. Traina F, De Fine M, Abati CN, Bordini B, Toni A. *Arch Orthop Trauma Surg*. 2012 Aug;132(8):1133-9. doi: 10.1007/s00402-012-1538-z. Epub 2012 May 10.

Abductor muscle strengthening in THA patients operated with minimally invasive anterolateral approach for developmental hip dysplasia. Benedetti MG, Cavazzuti L, Amabile M, Tassinari E, Valente G, Zanotti G, Vaianti E, Orsini S, Mariani E, Taddei F. *Hip Int*. 2019 Sep 22;1120700019877174. doi: 10.1177/1120700019877174.

A reproducible and inexpensive method of measuring hip abductor strength. Traina F, Cristofolini L, De Fine M, Brunetti L, Frakulli I, Abati CN, D'Uva L, Toni A. *Hip Int*. 2010 Oct-Dec;20(4):512-7. Doi: 10.1177/112070001002000415.

Long-term results of total hip replacement in patients with Legg-Calvé-Perthes disease. Traina F, De Fine M, Sudanese A, Calderoni PP, Tassinari E, Toni A. *J Bone Joint Surg Am*. 2011 Apr 6;93(7):e25. doi: 10.2106/JBJS.J.00648.

Outcome of hybrid stem fixation in osteoporotic female patients. A minimum five-year follow-up study. Traina F, Tassinari E, Pilla F, De Fine M, Biondi F, Cristofolini L, Toni A. *Int Orthop*. 2009 Dec;33(6):1489-94. doi: 10.1007/s00264-008-0645-5. Epub 2008 Aug 20.

Is a minimally invasive anterior approach effective in old patients? A pilot study. Faldini C, Perna F, Pilla F, Stefanini N, Pungetti C, Persiani V, Traina F. *J Biol Regul Homeost Agents*. 2016 Oct-Dec;30(4 Suppl 1):193-199

Total hip arthroplasty and sexual activity: a systematic review. Neonakis EM, Perna F, Traina F, Faldini O, Antoniou G, Kyriakopoulos G, Triantafyllopoulos IK, Megaloikononimos PD, Faldini C. *Musculoskelet Surg*. 2020 Apr;104(1):17-24. doi: 10.1007/s12306-020-00645-z. Epub 2020 Feb 6.

Highly Porous Titanium Cups versus Hydroxyapatite-Coated Sockets: Midterm Results in Metachronous Bilateral Total Hip Arthroplasty. Castagnini F, Bordini B, Yorifuji M, Giardina F, Natali S, Pardo F, Traina F. *Med Princ Pract*. 2019;28(6):559-565. doi: 10.1159/000500876. Epub 2019 May 13.

Alumina-on-alumina hip implants: A wear study of retrieved components. Affatato S, Traina F, De Fine M, Carmignato S, Toni A. *J Bone Joint Surg Br.* 2012 Jan;94(1):37-42. doi: 10.1302/0301-620X.94B1.26956.

Ceramic-on-Ceramic Total Hip Arthroplasty with Large Diameter Heads: A Systematic Review. Castagnini F, Cosentino M, Bracci G, Masetti C, Faldini C, Traina F. *Med Princ Pract.* 2020 Aug 5:1-8. doi: 10.1159/000508982.

Mixed ceramic combinations in primary total hip arthroplasty achieved reassuring mid-to-longterm outcomes. Castagnini F, Bordini B, Biondi F, Cosentino M, Stea S, Traina F. *J Mater Sci Mater Med.* 2020 Jun 27;31(7):56. doi: 10.1007/s10856-020-06393-7.

Effect of cup medialization on primary stability of press-fit acetabular cups. Morosato F, Cristofolini L, Castagnini F, Traina F. *Clin Biomech (Bristol, Avon).* 2020 Sep 7;80:105172. doi: 10.1016/j.clinbiomech.2020.105172.

Patient weight more than body mass index influences total hip arthroplasty long term survival. Traina F, Bordini B, De Fine M, Toni A. *Hip Int.* 2011 Nov-Dec;21(6):694-9. doi: 10.5301/HIP.2011.8879.